



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The osmoresponsiveness of oxytocin and vasopressin neurones: mechanisms, allostasis and evolution

Citation for published version:

Leng, G & Russell, JA 2018, 'The osmoresponsiveness of oxytocin and vasopressin neurones: mechanisms, allostasis and evolution', *Journal of Neuroendocrinology*. <https://doi.org/10.1111/jne.12662>

Digital Object Identifier (DOI):

[10.1111/jne.12662](https://doi.org/10.1111/jne.12662)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Neuroendocrinology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Journal of Neuroendocrinology

The osmore sponsiveness of oxytocin and vasopressin neurones: mechanisms, allostasis and evolution.

Journal:	<i>Journal of Neuroendocrinology</i>
Manuscript ID	JNE-18-0124-RA.R1
Manuscript Type:	Review Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Leng, Gareth; University of Edinburgh, Centre for Discovery Brain Sciences Russell, John
Keywords:	Vasopressin, Oxytocin, osmoreceptors, supraoptic nucleus

SCHOLARONE™
Manuscripts

The osmosensitiveness of oxytocin and vasopressin neurones: mechanisms, allostasis and evolution.

Gareth Leng and John A. Russell

Centre for Discovery Brain Sciences, The University of Edinburgh UK

Abstract

In the rat supraoptic nucleus, every oxytocin cell projects to the posterior pituitary, and is involved in both reflex milk ejection during lactation, and in regulating uterine contractions during parturition. All are also osmosensitive, regulating natriuresis. All are also regulated by signals that control appetite, including neural and hormonal signals that arise from the gut after food intake and from the sites of energy storage. All are also involved in sexual behaviour, anxiety-related behaviours, and social behaviours. The challenge is to understand how a single population of neurones can coherently regulate such a diverse set of functions, and adapt to changing physiological states. Their multiple functions arise from complex intrinsic properties which confer sensitivity to a wide range of internal and environmental signals. Many of these properties have a distant evolutionary origin, in multi-functional, multisensory neurones of *Urbilateria*, the hypothesised common ancestor of vertebrates, insects and worms. Their properties allow different patterns of oxytocin release into the circulation from their axon terminals in the posterior pituitary, into other brain areas from axonal projections, and independent release from their dendrites.

Introduction

In 1989, in the first issue of this journal, we, with Richard Dyball and Ruth Blackburn, published a paper entitled “*Role of anterior peri-third ventricular structures in the regulation of supraoptic neuronal activity and neurohypophysial secretion in the rat*” (1). Despite the less than catchy title, the *Web of Science* records that it has been cited 121 times. The studies it describes resolved an argument between us, and the notice that it received is an indication that there were many parties to that argument. So what exactly were the controversial issues?

Osmoregulation and neurohypophysial hormones

1
2
3 In the 1940's Verney (2) had established that, when water and salt balance are threatened by
4 dehydration, 'osmoreceptors' in the brain detected such threats and mediated stimulation of
5 vasopressin secretion from the posterior pituitary gland to counteract the disturbance. He
6 suggested that these were in the supraoptic nucleus of the hypothalamus, functioning as "stretch
7 receptors" that directly stimulated the neurones of the hypothalamo-hypophysial tract.
8
9

10
11 Our 1989 paper addressed the issue of the osmosensitiveness of magnocellular
12 neurones, in terms of whether these neurones are directly osmosensitive, or whether they respond
13 to inputs from osmoreceptors elsewhere. To understand the argument it is germane to note that
14 one of us (GL) was an electrophysiologist, working in Barry Cross's group at Babraham, while
15 the other (JAR) was a classical physiologist, mentored by Mary Pickford, at Edinburgh. Pickford
16 had pioneered our understanding of how afferent signals regulated these neurones, particularly
17 by her studies of the regulation of antidiuresis by central acetylcholine (3).
18
19

20
21 *Distinguishing oxytocin and vasopressin cells.* In 1959, Cross and Green made the first
22 attempt to study the electrical behaviour of magnocellular neurones in response to hyperosmotic
23 stimulation (4), but this was confounded by the difficulty in distinguishing them from
24 neighbouring neurones. However, the introduction of *antidromic identification* in the 1970s
25 enabled electrophysiological recordings to be made from neurones identified as projecting to the
26 posterior pituitary gland. This technique involves placing a stimulating electrode on the neural
27 stalk by which neurones that project to the posterior pituitary can be positively identified by the
28 appearance of fixed latency action potentials evoked "antidromically" by stimuli applied to the
29 stalk. It soon became apparent that oxytocin cells as well as vasopressin cells responded to
30 osmotic stimulation. In response to systemic osmotic stimulation, oxytocin cells fired
31 continuously at an elevated rate, while many vasopressin cells fired phasically, in long bursts of
32 spikes, with the duration of bursts and the spike frequency within bursts determining the rate of
33 vasopressin secretion (5).
34
35

36
37 *For direct osmosensitivity.* By 1989, there was compelling electrophysiological evidence
38 that magnocellular neurones were directly osmosensitive; direct application of hypertonic saline
39 would excite these cells *in vivo*, (6) and the first intracellular recordings *in vitro* had shown that
40 they were depolarized by an increase in extracellular osmotic pressure even when all synaptic
41 input was blocked (7). However, classical physiologists remained sceptical: they noted that the
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

experimental conditions of the *in vitro* electrophysiological studies were ‘unphysiological’, and that the saline doses applied were excessively (“supraphysiologically”) high.

For osmosensitive inputs. Just as compelling to physiologists as the electrophysiological evidence was to electrophysiologists was evidence that lesions of brain regions anterior to the supraoptic nucleus eliminated the osmotic regulation of vasopressin secretion (8-11). Typically, these lesions encompassed the ventral median preoptic nucleus (also called the nucleus medianus), the organum vasculosum of the lamina terminalis (OVLT), and periventricular tissue caudal and lateral to the ventral lamina terminalis, and produced retrograde degeneration in the subfornical organ and terminal degeneration in the supraoptic nucleus (12). Lesions to the subfornical organ alone also produced a reduction on osmotically-evoked vasopressin release (13).

This focused attention on three sites: two of these, the subfornical organ and OVLT (14) are densely vascularized circumventricular organs and lie outside the blood-brain barrier, apparently ideally situated to monitor the composition of the blood. The third site, the nucleus medianus, lies between the subfornical organ and the OVLT and receives a dense synaptic input from both (15). Collectively these came to be known as the AV3V region – the region anterior and ventral to the third ventricle. All three sites projected densely to the magnocellular neurons (15).

Recalling those past clashes, several themes stand out. The evidence itself was not in dispute: both ‘sides’ of the evidence had been replicated extensively. Accordingly, the challenge was to construct a credible narrative that accommodated all the evidence. There were two important impediments: skepticism about how the same neuroendocrine system could simultaneously regulate such disparate functions as electrolyte homeostasis and the reproductive functions, and the counter-intuitive notion that neurones might be osmosensitive despite evidence that their osmosensiveness was selectively eliminated by lesions of an afferent pathway.

The osmosensiveness of magnocellular neurones.

Evidence for the importance of circumventricular organs came from studies that addressed three things of physiological importance: thirst, sodium excretion, and urine production. Each of these was profoundly impaired by lesions to any part of the AV3V region.

1
2
3 The effects on urine flow clearly reflected a loss of the antidiuretic actions of vasopressin, while
4 effects on thirst were believed to reflect actions independent of pituitary hormone secretion (16).
5 However, the effects on sodium excretion were problematic. In some species, vasopressin
6 contributes to sodium excretion (17), but this could not account for the observed deficit – there
7 was a missing “natriuretic factor”: *“It is thus probable that a cerebral natriuretic system is*
8 *involved in the functional expression of any other peripheral natriuretic system, e.g. the heart*
9 *atrial natriuretic system”* (18).

15 A possibility was that, in at least some species, oxytocin might be such a natriuretic
16 factor, as Brooks and Pickford had shown in the dog that oxytocin could increase sodium
17 excretion (3), and oxytocin also had natriuretic actions in the rat (19-21). The
18 electrophysiological studies implied that, in the rat, oxytocin cells were just as osmosensitive as
19 vasopressin cells, and Young and van Dyke had in 1968 shown that in the rat progressive
20 dehydration reduced the neurohypophysial content of both oxytocin and vasopressin to a similar
21 extent (22). However, oxytocin was the hormone of milk-ejection and parturition, and a
22 predominant assumption was that different physiological functions were compartmented in
23 different neuronal populations – and that different hormones had separate physiological roles.
24 There appeared to be no osmotic regulation of oxytocin secretion in humans (23, 24) and no
25 clear renal actions (25). Thus in 1974, Lee and de Wardener declared: *“One cannot better the*
26 *conclusions reached by Bentley in 1971 (26) that in mammals the neurohypophysial hormones*
27 *may, in an unpredictable way, increase sodium excretion in the rat, dog, camel and sheep, but*
28 *not man”* (27).

39 However, by 1989, the osmoresponsiveness of oxytocin cells in rats had been shown
40 from many studies. Dehydration and sodium loading by intraperitoneal injection of hypertonic
41 saline increased the electrical activity of oxytocin cells *in vivo* and increased oxytocin secretion
42 as strongly as they increased the activity of vasopressin cells and vasopressin secretion (28, 29).
43 Moreover, in rats, oxytocin had natriuretic effects at low doses (24, 30), and evidence
44 accumulated that osmotically-stimulated oxytocin secretion contributed to natriuresis (31, 32)
45 both by possible direct actions at the kidney (24) and by regulating the secretion of atrial
46 natriuretic peptide from the heart (33). Lesions to the AV3V region abolished not only
47 osmotically-induced vasopressin secretion (34), but also osmotically-induced oxytocin secretion,
48 and blocked increased synthesis of both peptides in response to water deprivation (8). By
49
50
51
52
53
54
55
56
57
58
59
60

contrast, AV3V lesions did so without affecting suckling-induced oxytocin release (35) or parturition (36)— the progress of which, in rats, is driven by uterine contractions that activate oxytocin cells via an input from the caudal brainstem (37-39).

Intrinsic osmosensitivity dependence on an excitatory input. To reconcile the electrophysiological evidence with the results of lesion studies, it was necessary to explain how a lesion to an afferent input selectively impaired the ability of magnocellular neurones to express their intrinsic osmosensitivity. That explanation, as first presented (6, 40) acknowledged that the spiking activity of magnocellular neurones depends on excitatory synaptic inputs, but proposed that the frequency at which spikes occurred in response to such inputs depends on the level of intrinsic depolarization. Accordingly, an input might be essential for osmoresponsiveness even if it was not itself osmoregulated. This notion, that neuronal “noise” might be important, was a seeming affront to the idea that spike activity in neurones was the harbour of physiologically meaningful information in the brain.

It was this hypothesis, disputed amongst us, that we put to collaborative test in 1989 (1). In anesthetized rats, we lesioned the AV3V region rendering the magnocellular neurones silent and unresponsive to systemic osmotic stimulation. We then restored normal levels of electrical activity by continuous ejection of glutamate from the recording microelectrode. We found that this rescued the ability of the neurones to respond to systemic osmotic stimulation, showing that their intrinsic osmosensitivity was indeed sufficient to modulate their firing rate in the presence of an input that was not itself osmoresponsive. However, the extent of the activation was less than in normal rats, indicating that normal osmoresponsiveness involves both intrinsic osmosensitivity and increased synaptic input.

The direct osmosensitive mechanism can depolarize magnocellular neurones by only a few millivolts—too little to explain, on its own, the changes in spike activity that physiological increases in plasma osmotic pressure produce (Figure 1).

However, if the membrane potential of a neurone is continually fluctuating, a small sustained depolarization, by altering the probability that fluctuations will exceed the spike threshold, will increase the firing rate. This phenomenon by which noise enhances the sensitivity of neurones is called *stochastic resonance* and is now recognized as a general feature of sensory systems (43).

Osmosensitive mechanisms. The osmosensitivity of magnocellular neurones involves specialised stretch-sensitive ion channels, as shown by Oliet and Bourque in 1993. When the

1
2
3 extracellular osmotic pressure rises, the cells shrink, and this opens stretch-sensitive membrane
4 ion channels causing a depolarizing current to flow (44). This involves an N-terminal variant of
5 the transient receptor potential vanilloid 1 (Trpv1) channel, activation of which triggers a
6 mechanical process that engages a thin layer of actin filaments (F-actin) beneath the plasma
7 membrane, and a network of microtubules (45, 46). The same mechanism contributes to the
8 osmotic regulation of thirst by neurones within the AV3V region (47).
9

10
11
12
13 The experiments implicating this channel involved local application of mannitol as a
14 hypertonic stimulus. However, mice lacking the Trpv1 channel show normal vasopressin
15 secretion and normal thirst in response to hypernatremia (48), raising the possibility that the
16 Trpv1 channel contributes to vasopressin secretion and thirst stimulated by hyperosmolality but
17 not that stimulated by hypernatremia. Consistent with this, Kinsman (49) reported that systemic
18 injection of mannitol stimulates thirst similarly in normal mice and Trpv1 knockout mice. Thus
19 magnocellular neurones are directly sensitive to both osmotic pressure and sodium.
20
21

22
23
24
25 Magnocellular neurones express two other members of the Trpv family of channels. In the
26 paraventricular nucleus, Trpv4 is expressed selectively in magnocellular vasopressin cells (50),
27 and seems also to be involved in osmoresponsiveness (51, 52). Trpv2 is also expressed densely
28 in the supraoptic nucleus and the magnocellular portion of the paraventricular nucleus in both
29 oxytocin cells and vasopressin cells (53, 54); little is known of its function in these cells, but in
30 other tissues Trpv2 channels have been associated with mechanosensitivity, thermosensitivity
31 and osmosensitivity (51). The Trpv1 channels that mediate osmosensitivity also confer
32 thermosensitivity on the vasopressin cells (55, 56): vasopressin is released in hot conditions to
33 preserve body water in the face of evaporative loss.
34
35

36
37
38
39
40
41 *Degeneracy.* Osmoresponsiveness thus involves multiple 'degenerate' mechanisms.
42
43 *Degeneracy* refers to different mechanisms that converge to produce the same result, whereas
44 *redundancy* refers to duplication of a mechanism (57). Degeneracy contributes to robustness in
45 biological systems, and it has been argued that the evolution of degenerate mechanisms is a
46 common consequence of natural selection, as there is little selection pressure for the elimination
47 of either neutral or degenerate mutations. Several types of sodium channel appear to contribute
48 to sodium detection in the supraoptic nucleus (58-60). Osmotic stimuli also promote the
49 phosphorylation of extracellular signal-regulated protein kinases in magnocellular neurones and
50
51
52
53
54
55
56
57
58
59
60

in many neurones of the AV3V region, and this modulates osmotransduction by a mechanism still undetermined (61).

Complex osmosensitive inputs. The osmosensiveness of magnocellular neurones involves many other factors (62), including afferent signals from the AV3V region; some of which are also osmosensitive. The OVLT and subfornical organ also contain neurones that respond to blood-borne hormones that have an important role in electrolyte homeostasis, including angiotensin II, relaxin and atrial natriuretic peptide (63). As well as conventional neurotransmitters, efferent signals from the AV3V region involve a variety of peptides including angiotensin II from the subfornical organ (64). Angiotensin II has opposite effects on vasopressin and oxytocin cells: in both it opens the channels that mediate osmosensitivity (65, 66), but in oxytocin cells it also induces endocannabinoid release, which opposes the excitatory effects of OVLT stimulation. Thus angiotensin promotes antidiuresis but inhibits natriuresis, as seems appropriate for a signal primarily regulating blood volume and activated by hypovolemia.

Role for glia. The osmosensiveness of supraoptic neurones is modulated by taurine from local astrocytes; taurine is an osmolyte which is actively exported from many cells under hypotonic conditions to help maintain cell volume homeostasis, and hypertonic aCSF microdialysed into the supraoptic nucleus strongly stimulates Fos and *c-fos* mRNA expression in astroglia in this nucleus (67). Taurine is an agonist at glycine receptors; these ligand-gated chloride channels are expressed in supraoptic neurones, and because the electrochemical gradient for chloride favors influx under resting conditions, taurine promotes hyperpolarization, moderating the gain of their excitatory response to hypertonic stimuli (68, 69).

Importance of inhibitory input. The AV3V region provides both excitatory and inhibitory synaptic inputs to magnocellular neurones (70). The osmoreceptive neurones in the OVLT that project to the supraoptic nucleus are all glutamatergic (71), but the OVLT and the subfornical organ also project indirectly via the median preoptic nucleus – and this involves the inhibitory transmitter GABA (72). Systemic osmotic stimulation thus triggers the release of both GABA and glutamate in the supraoptic nucleus, as confirmed by microdialysis *in vivo* (41). While it might seem perverse that magnocellular neurones receive an osmotically regulated input that comprises a mixture of inhibition and excitation, this might be adaptive. The response of vasopressin secretion to increasing osmotic pressure is linear over the range of osmotic pressure experienced by animals over prolonged water deprivation, and linearity is also apparent in the

1
2
3 responses of individual oxytocin cells and vasopressin cells *in vivo*. This linearity is surprising:
4 neurones typically become more excitable as they are excited— an EPSP is more likely to cross
5 the spike threshold when a neurone is partially depolarised, so their response to an increasing
6 excitatory input tends to increase non-linearly. This non-linearity truncates their dynamic range,
7 as they reach maximum firing rates more quickly. However, we noted that if an input comprised
8 a mixture of synaptic excitation and synaptic inhibition then the neuronal response to increasing
9 input rates would be more linear and the dynamic range would be extended (73). It might be
10 expected that an excitatory input would be cancelled out by an equal and opposite inhibitory
11 input, but this is not the case for random inputs. A mixed random input produces a membrane
12 potential that fluctuates around the mean; spikes arise when fluctuations exceed spike threshold,
13 and these spike triggering events increase in frequency linearly with the mean input rate (41, 73).
14
15

16
17 In normal rats, GABA inhibits both oxytocin and vasopressin cells *in vivo*. As mentioned,
18 stimulation of the OVLT region produces mixed excitatory and inhibitory effects on oxytocin
19 and vasopressin cells *in vivo*, with the inhibitory effects arising via activation of a GABAergic
20 input from the nucleus medianus. This inhibitory effect can be blocked by microdialysis of the
21 GABA antagonist bicuculline onto the supraoptic nucleus (74) (Figure 2), as can inhibition
22 arising from stimulation of the arcuate nucleus (75).
23

24
25 In magnocellular vasopressin cells in slice preparations *in vitro*, excitatory responses to
26 GABA have been reported in some experimental conditions (76, 77) though not in others (78).
27 This indicates that the intracellular chloride concentration, which determines the direction of the
28 neuronal response to GABA, is vulnerable to particular experimental conditions, and raises the
29 question of whether a similar change occurs in physiological conditions. This might be
30 anticipated in conditions of chronically sustained activation of GABA inputs which lead to a
31 sustained elevation of chloride entry. Systemic osmotic stimulation, as indicated above, involves
32 activation of GABA inputs to the magnocellular neurones, and chronic salt loading indeed leads
33 to a change in the direction of GABA actions (78). Two other studies have indicated that the
34 direction of GABA actions can change from inhibition to excitation in conditions of chronic
35 hyperactivation of vasopressin secretion (79, 80), and one has reported a change affecting both
36 oxytocin and vasopressin cells in lactation (81).
37
38

39
40 *Wider inferences.* These issues were harbingers of the coming revolution in our
41 understanding not just of the magnocellular neurones but of neuroendocrine systems in general.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Today, it is accepted that neuroendocrine neurones are multifunctional; that they express a wide range of properties that make them directly sensitive to their immediate external environment (Figure 3); and that they are phenotypically plastic, with properties that vary with physiological state.

It is also now clear that both populations of magnocellular neurones are heterogeneous in their intrinsic properties. These neurones have some features that distinguish them clearly from most other hypothalamic neurones, and some features that are more commonly observed in vasopressin cells than in oxytocin cells and vice versa, but there is considerable variation within each of these populations (85). A few cells appear to have an ambiguous phenotype – for example a few cells generate both suckling-induced milk-ejection bursts, classically identifying them as oxytocin cells, but also show phasic firing patterns that are mainly associated with vasopressin cells. Most magnocellular neurones express mRNAs for both oxytocin and vasopressin; usually these are present at very different levels but a small proportion (~3%) express both at equivalent levels. Interestingly, in conditions of sustained elevated demand, the proportion that express appreciable amounts of both peptides increases (to 24% in the study of da Silva *et al.* (86)).

Multi-functional magnocellular neurones

Different physiological responses arise in part from different patterns of activity. In response to raised plasma osmotic pressure, oxytocin cells fire continuously (41), but in response to suckling and during parturition they fire in intense synchronised bursts every few minutes. These bursts lead to a secretion that is amplified by non-linearities in stimulus-secretion coupling at the nerve terminals (87), resulting in a sequence of large pulses of oxytocin secretion. Only at term pregnancy does the uterus express abundant receptors for oxytocin, and only in lactation does the mammary gland. The mammary gland “senses” only pulses of secretion: being relatively insensitive to oxytocin, the mammary gland is indifferent to the lower concentrations induced by osmotic challenge. By contrast, the kidney responds to secretion evoked by small, sustained increases in oxytocin secretion, and is indifferent to brief intermittent pulses (88, 89). Thus oxytocin cells can regulate milk-let down and natriuresis simultaneously without conflict.

Salt appetite. Electrolyte homeostasis necessarily involves regulation of both sodium excretion and sodium intake. Dietary sodium deprivation elicits a strong salt appetite in rats (90), as does bilateral adrenalectomy (91, 92); aldosterone through its actions on the brain (93, 94);

and hypovolemic stimuli (95-97). The AV3V region is strongly implicated in salt appetite, in part through central angiotensin II pathways from the subfornical organ (98-102).

However, sodium intake is not always stimulated when hypovolemia is present or when blood angiotensin II levels are elevated. Diverse treatments inhibit salt appetite, including acute hyperosmolality, uremia, severe hypovolemia, hypotension and nausea (103) – all stimuli that increase oxytocin secretion. Conversely, angiotensin-II-induced salt intake is potentiated by treatments that decrease oxytocin secretion, including systemic injection of deoxycorticosterone (104) or ethanol (105), and is also potentiated by destruction of central neurones bearing oxytocin receptors (106) (107) and by pretreatment with an oxytocin antagonist (108). In rats, sodium depletion evokes a powerful and selective sodium appetite, and many studies have shown that centrally administered oxytocin inhibits this, as do many physiological stimuli that increase oxytocin secretion, such as dehydration or salt loading. Naloxone, which augments stimulated oxytocin secretion, inhibits salt appetite induced by colloid treatment, and this is abolished by i.c.v. pretreatment with an oxytocin receptor antagonist (103). It has also been proposed that the neuropeptide adrenomedullin inhibits salt appetite via its effects on oxytocin release (109). Thus there is extensive evidence that, in rats, central release of oxytocin suppresses salt appetite: an effect complementing its peripheral natriuretic action.

The sites at which oxytocin modulates salt appetite may include the AV3V region itself, as the OVLT contains oxytocin fibres (104), and there is electrophysiological evidence that some magnocellular neurones of the supraoptic nucleus project to that region (110). Another key site is the parabrachial nucleus, where oxytocin receptor-expressing neurones have been implicated in the regulation of water and saline intake (111, 112).

Food intake. Salt appetite is a conspicuous feature of animals whose diet is mainly vegetarian, and which may accordingly have difficulty in gaining enough sodium in their diet to meet their needs. Humans, by contrast, do not exhibit a strong salt appetite even in conditions of hyponatremia. However, oxytocin is involved in the regulation of food intake more generally. Both oxytocin cells and vasopressin cells are activated acutely by food ingestion (113, 114). This might be an anticipatory response to the electrolyte imbalance that will arise from solute intake, but oxytocin has a now well-established central role in energy balance: in rats it suppresses voluntary intake of sweet carbohydrates, and in mice it promotes energy expenditure and thermogenesis (115). The oxytocin cells of the rat supraoptic nucleus express insulin receptors

and glucokinase, and are activated by both glucose and insulin (116). They also express receptors for leptin and insulin, as well as for many anorectic peptides released from the brain itself, such as α -MSH (melanocyte stimulating hormone) (117), and they are activated by systemic administration of leptin, secretin and cholecystokinin (CCK) (115, 118, 119). The effects of CCK (120) and probably those of secretin too are mediated by activation of gastric vagal afferents that lead to activation of neurones in the nucleus tractus solitarii (NTS) (121), including the noradrenergic A2 cells that project directly to oxytocin cells. Oxytocin cells are also activated during voluntary food intake (113), by intragastric gavage of the dietary amino acid L-tryptophan (122), and by gavage of sweet energy dense food, but interestingly they are inhibited by gavage of isocaloric cream (123). Many secreted peptides are co-expressed with oxytocin and/or vasopressin, and some of these, like nesfatin (124, 125) CCK (126), galanin-like peptide (127) and pituitary adenylate cyclase activating polypeptide (PACAP) (128) have anorectic effects that may support the anorectic effects of oxytocin at central sites. The central sites at which oxytocin exerts its effects on feeding are likely to include the ventromedial nucleus of the hypothalamus (115) and the central amygdala, two sites that express oxytocin receptors densely; the central amygdala is involved in salt appetite (129) as well as more generally in food intake (130) and receives a projection from magnocellular oxytocin cells (131). Thus the roles of central oxytocin on appetite go far beyond the regulation of salt intake.

Allostasis.

The well-established principle of physiology is that *homeostasis* maintains a constant internal environment, and theoretical models have envisaged a ‘set-point’ that a control system aims at maintaining. However, controlled variables vary with functional demand and environment, while control mechanisms are expected to settle at a level that uses least energy to resolve challenges- i.e. *allostasis*, or stability through change, including in anticipation of future demands (132). Anticipatory changes in vasopressin secretion and thirst occur in response to water intake in advance of any change in plasma $[Na^+]$ (133, 134), thirst (135) is activated independently of plasma $[Na^+]$ by circadian cues mediated by a projection from the suprachiasmatic nucleus to the OVLT, and the osmoresponsiveness of magnocellular neurones is also modulated by a circadian input (136).

During chronic dehydration or salt loading there are extensive changes to the magnocellular system, as apparent from analyses of the transcriptome of the rat supraoptic nucleus (137-139). The changes include hypertrophy of the magnocellular neurones, increased synthesis of their products, intracellular machinery and various transcription factors, a re-organisation of the neurone-glia architecture (140), and increased expression of the gaseous transmitters nitric oxide and carbon monoxide (141).

Allostasis in pregnancy. Allostatic adaptations also accompany pregnancy. While normal homeostatic mechanisms serve to maintain a constant blood volume, a constant electrolyte composition of the blood and a stable body weight, pregnancy requires an expanded blood volume to support the metabolic demands of the growing fetus and an expansion of fat mass in preparation for meeting the nutritional demands of the newborn (142). To accommodate these requires a re-setting of these homeostatic set points, and this involves a complex array of adaptations that affect the oxytocin cells (137, 143-146).

Resetting of the set-points for volume and electrolyte balance arises in part from the actions on the subfornical organ and OVLT of *relaxin*, a peptide hormone produced by corpora lutea in pregnancy in increasing amounts as pregnancy progresses. Circulating relaxin stimulates both water intake (147) and vasopressin secretion via its actions on the AV3V region, and the combination of increased water intake and increased water retention contribute to a dilutional expansion of plasma volume with accompanying hyponatremia (148). This reduces plasma osmolality to below the normal set point for osmotic stimulation of oxytocin and vasopressin secretion (149). However, a reduction in the activity of oxytocin cells would entail a reduction in oxytocin synthesis, as observed with experimentally induced hyponatremia – and this would not be desirable, as the pituitary stores of oxytocin need to be expanded in preparation for the demands of parturition and subsequent lactation. However, relaxin also stimulates oxytocin neuronal activity (150), maintaining the normal level of synthesis. At the same time, oxytocin secretion needs to be restrained, both to minimise natriuresis and to help expand the pituitary store of oxytocin. This is achieved through another adaptation of pregnancy, involving the opioid peptide dynorphin.

Autoregulation of oxytocin secretion by dynorphin. Both magnocellular vasopressin cells and oxytocin cells co-express dynorphin, which acts at κ -opioid receptors on the cells of origin. In vasopressin cells, somato-dendritic release of dynorphin has a role in the phasic patterning of

spike activity (151), but in oxytocin cells dynorphin is an inhibitory feedback regulator of secretion from the pituitary, and upregulation of dynorphin expression in pregnancy contributes to the accumulation of pituitary oxytocin content in preparation for parturition (39). Excitation of the nerve terminals in the posterior pituitary releases dynorphin along with oxytocin, and this normally restrains activity-dependent secretion (152, 153). This effect is enhanced in early pregnancy, contributing to the accumulation of pituitary oxytocin content, but towards the end of pregnancy, it fades to leave action potentials more effective in stimulating oxytocin secretion (152).

In the pregnant rat, oxytocin cells will barely respond to modestly increased plasma $[Na^+]$; if the actions of dynorphin at the nerve terminals are blocked by the opioid antagonist naloxone, the response is enhanced, but is still weaker than in virgins (Figure 4). However, if plasma $[Na^+]$ is further increased, the response is greater in late pregnant rats than in virgins (154); hence, in late pregnancy plasma $[Na^+]$ is reduced below the threshold to stimulate oxytocin cells, but above this threshold the gain of their response is increased, probably as a reflection of the increased oxytocin store.

Central opioid mechanisms. Towards the end of pregnancy, the restraining influence of dynorphin wanes, but the activity of oxytocin cells must still be kept in check until the birth canal is ready for parturition. This check is effected by inhibition of oxytocin cell activity by opioid peptides that act at μ -opioid receptors. This involves opioid actions on afferent inputs to the oxytocin cells from the caudal brainstem (155), and might also involve direct actions, possibly by an input from arcuate nucleus β -endorphin neurones (156).

Increased food intake is an appropriate adaptation in late pregnancy, The mechanisms are complex (157), but decreased stimulation of oxytocin cells by appetite-related signals might be a contributing factor (158). Oxytocin cells are directly innervated by A2 noradrenergic neurones of the NTS, and this pathway is activated by gastric distension and by systemic administration of the gut hormone cholecystokinin (CCK). The A2 projection also carries inflammatory signals, and information from the contracting uterus (37), though whether these are conveyed through the same A2 neurones or different subsets is not known. Certainly the A2 cell group as a whole is functionally heterogeneous, as CCK preferentially activates Fos expression in the subset that projects to the supraoptic nucleus (120). The noradrenergic nerve terminals are regulated presynaptically by an opioid peptide that acts at μ -opioid receptors (159). As A2 neurones co-

express both pro-enkephalin-A and μ -opioid receptor mRNAs (160), it seems likely that this reflects an autoregulatory brake on noradrenaline release analogous to the dynorphin brake on oxytocin secretion from the pituitary. In late pregnancy there is increased expression of both pro-enkephalin-A and μ -opioid receptor mRNAs in the NTS (160), and at this time (but not in virgin rats or in lactation (161)), naloxone enhances the activation of oxytocin cells by CCK (155). Thus, in late pregnancy, there is enhanced opioid restraint of the noradrenergic projection. We measured oxytocin secretion in response to different stimuli in virgin and late pregnant rats, and in these experiments we blocked all opioid actions by pretreating the rats with naloxone (Figure 4). Both AV3V stimulation and i.p. hypertonic saline stimulated oxytocin secretion less effectively in late pregnant rats, seeming to suggest a reduction in the drive by AV3V inputs, while oxytocin secretion in response to CCK or interleukin-1 β (IL-1 β) was greater, suggesting that the brainstem input to the oxytocin cells is more effective. However, the reduced response to an osmotic challenge at least in part reflects the chronic hyponatremia that is present in late pregnancy, and this might also affect the response to AV3V stimulation. Moreover, the enhanced response to CCK and IL-1 β in part reflect the increase in the availability of oxytocin for activity dependent release that results from the accumulation of pituitary content that occurs in pregnancy. What we see from such experiments is that pregnancy entails multiple changes in the oxytocin system, changes in the oxytocin cells themselves, in their inputs, and in stimulus – secretion coupling at the nerve terminals; some of these alter the responsiveness of oxytocin secretion to particular input(s), but other changes are necessary to maintain normal responsiveness.

Allopregnanolone. In considering what might drive the expression of pro-enkephalin-A mRNA in the NTS in pregnancy, we ruled out progesterone, having found very few neurones expressing the progesterone receptor in the NTS (162). However, the high levels of progesterone in pregnancy are associated with increased levels of its metabolite allopregnanolone in the circulation and in the brain. Allopregnanolone is an allosteric modulator of GABA_A receptors, prolonging their opening time when activated by GABA, and in late pregnancy this also enhances GABAergic inhibition of oxytocin cells (163, 164). Expression of mRNA for 5 α -reductase, the rate-limiting enzyme in allopregnanolone production, is increased in late pregnancy, and blocking this enzyme with finasteride reduces pro-enkephalin-A expression in the NTS (165).

The consequences of this can be seen by studying oxytocin release in response to systemic administration of IL-1 β ; this increases the firing rate of oxytocin cells in virgin rats, but not in late pregnant rats unless naloxone is given just before IL-1 β (166). Similarly IL-1 β increases Fos expression in virgin but not pregnant rats, and again the responsiveness to IL-1 β is rescued by pre-treatment with naloxone, or by blocking allopregnanolone production with finasteride. Conversely allopregnanolone treatment of virgin rats suppresses the responsiveness of oxytocin neurones to IL-1 β (167).

Allopregnanolone also moderates the functional effect of oxytocin in the spinal cord. The spinal cord receives a dense projection from a small population of oxytocin neurones in the paraventricular nucleus (168), and this pathway modulates pain sensitivity. Oxytocin-induced analgesia in the spinal cord appears to be mediated in part at least by activation of GABA release, and its effectiveness is amplified by the actions of allopregnanolone (169).

The evolution of magnocellular neurones

Oxytocin and vasopressin arose by duplication of the vasotocin gene at around the time of appearance of the earliest vertebrates. Most modern vertebrates have at least two oxytocin- and vasopressin-like peptides, while most invertebrates – including mollusks, annelids and many insects (170, 171) - have only one.

Fishes. The earliest vertebrate fossils are of jawless fish, like modern lampreys, and lampreys have only one peptide that is like oxytocin and vasopressin, vasotocin. Jawed fish, a category that includes cartilaginous fish and bony fish, appeared about 440 million years ago and all living jawed fishes have vasotocin and an oxytocin-like peptide. Bony fish all have vasotocin and isotocin, and include ray-finned fish and lobe-finned fish (such as lungfish). Amphibians, mammals, reptiles, and birds evolved from lobe-finned fish and all have at least one homolog of oxytocin and one of vasopressin.

In ray-finned fishes, vasotocin is involved in osmoregulation and isotocin in regulating electrolyte concentration, so it appears that, in the aquatic vertebrates where vasotocin and isotocin first appeared as distinct hormones, both were involved in water and electrolyte balance; they may also have been involved in reproduction, the timing of which is generally tied to environmental conditions. In bluehead wrasse, vasotocin influences social behavior and is regulated by sex steroids: (172). In medaka, isotocin expression is up-regulated selectively in

male brains by gonadal androgens (173, 174). In goldfish, isotocin is a regulator of food intake (175).

Out of water. In land vertebrates, isotocin evolved into mesotocin by a single amino acid substitution and from mesotocin into oxytocin by another. Vasotocin evolved into vasopressin; again with a single amino acid substitution. Thus vasopressin and oxytocin arose through duplication of the vasotocin gene in a species of jawless fish that lived about 400 million years ago. When a gene is duplicated, one copy can maintain the original function, leaving the other free to mutate, and diverge, as happened here.

Receptors. Pituitary or peripherally released peptides cannot be anatomically confined — once released, they can diffuse or be conveyed by the blood, hemolymph, or extracellular fluid to distant sites; accordingly, for co-existing descendants of such a peptide to acquire differentiated functions, they must act at different receptors. The Cambrian explosion involved two episodes of whole genome duplication, and these separated V1a, V1b and oxytocin receptors; V2 receptors had apparently already separated from an ancestor of the V1 receptor family in an earlier gene duplication (176, 177).

Passwords for separate cell expression. When the vasotocin gene was duplicated, there were thus already two families of receptors present that could allow the functions of descendant peptides to diverge. However, for those functions to diverge, vasopressin and oxytocin had to be expressed in different cells. Every cell type has a molecular “password,” a combination of transcription factors that determine its identity, and genes with regulatory elements that recognize this password will be expressed in those cells (178). Murphy et al. (179) produced transgenic rats by inserting 40,000 bases of pufferfish DNA that included the isotocin gene. In these rats, isotocin was expressed only in oxytocin cells, and, in response to dehydration, expression of both isotocin and oxytocin were stimulated in a similar way. Thus mammalian oxytocin cells must have the same password as isotocin cells in fish, a password that arose early in vertebrate evolution and which has been conserved through subsequent evolution. Equally, the regulatory elements of oxytocin-like genes must also have appeared early in vertebrate evolution. In effect, the Cambrian explosion resulted in a duplication of the vasotocin cells, allowing these two sets of cells to diverge in function.

Conclusions

Oxytocin and vasopressin cells arose by duplication of a cassette of genes that defined a common neuronal phenotype, a phenotype that can be traced back to *Urbilateria*, hypothesized marine organisms that are proposed to be the last common ancestor of vertebrates, flies, and worms. In *Urbilateria*, cells that secreted a peptide ancestor of vasotocin apparently responded to diverse cues from their marine environment. These cells combined properties that we have thought of as separate properties of endocrine cells and neurons. They used a diversity of signaling mechanisms, made both peptides and neurotransmitters, and were endowed with a wide range of specialized senses, linking feeding, reproduction and internal homeostasis, to external conditions. They were not committed to a single role, but integrated multiple behavioral and physiological functions. In the nematode *C. elegans*, nematocin, a homolog of oxytocin and vasopressin, is critically involved in gustatory associative learning – the process by which nematode behaviour is modified by a learned association of salt and food availability (180).

Magnocellular neurones communicate with each other, with other neurones, and with other cells including glial cells. This communication involves many different messengers, including oxytocin and vasopressin but also other peptides including dynorphin that are co-packaged in the same vesicles as oxytocin and vasopressin, nitric oxide and prostaglandins that are produced *de novo*, and other neuromodulators such as adenosine (151). The release of these messengers is governed differentially; the rules that link neuronal activity to release vary according to the messenger concerned, and they differ between different sites of release within the same neurone, differ according to physiological state, and differ between oxytocin cells and vasopressin cells.

All magnocellular neurones express vesicle glutamate transporter-2 (181), indicating that they use glutamate as a conventional neurotransmitter at synaptic terminals. While all magnocellular neurones project to the posterior pituitary, subpopulations project to various central sites including the nucleus accumbens, the hippocampus, the amygdala, and the bed nucleus of the stria terminalis (182). Oxytocin receptors are expressed at these sites but also at many sites in the brain that receive few if any oxytocin fibres. It seems likely that within the brain, oxytocin cells release glutamate at synapses in the manner typical of neurotransmitters, but also release occasional vesicles containing oxytocin from axonal varicosities, to act as a local neuromodulator (183-188).

Vasopressin and oxytocin are released not only from nerve terminals in response to spike activity but also from the soma and dendrites in response to the mobilization of intracellular Ca^{2+} (189). Considerable amounts of peptide can be released from the dendrites; the oxytocin that is released from dendrites during suckling has a key role within the nucleus in generating milk-ejection bursts (190), but there is also considerable dendritic oxytocin release during parturition and sexual activity, and this appears to have a neurohormonal action at relatively distant sites to influence social behaviours (189, 191)

Like oxytocin, vasopressin has many behavioural effects. The role of magnocellular vasopressin cells is less well explored, but they are also active during feeding (114) and appear to have an anorectic action (192, 193). However, vasopressin is not only expressed in magnocellular neurones, but also in parvocellular neurones of the paraventricular nucleus that regulate the stress axis, in neurones of the suprachiasmatic nucleus that regulate circadian rhythms (194), and in diverse other populations, including in the olfactory bulbs (195) and retina (196). Accordingly, the behavioural roles of vasopressin might reflect compartmentalisation of function in different subsets of neurones. However, oxytocin cells do not allow this explanation: these are found *only* in the hypothalamus, and in the rat at least, only a few project exclusively within the brain (182), mainly to the dorsal vagal complex to regulate gastric reflexes and to the spinal cord to modulate penile erection and pain responses.

Thus we have to abandon any notion that oxytocin has a single function or even a single ‘main’ function. We must acknowledge that oxytocin cells are intrinsically multi-functional. Throughout their evolutionary history there has been co-evolution of the oxytocin system and its inputs to maintain that multi-functionality, and evolution of mechanisms that allow that multi-functionality to be adapted to meet changing physiological states. This means that we cannot interpret physiological changes in neuronal properties in isolation, but only in the context of everything else in the system that has changed. We cannot escape the necessity of building a systems level understanding to understand the importance of cellular properties.

References

1. Leng G, Blackburn RE, Dyball RE, Russell JA. Role of anterior peri-third ventricular structures in the regulation of supraoptic neuronal activity and neurohypophyseal hormone secretion in the rat. *J Neuroendocrinol.* 1989; **1**(1): 35-46.

2. Verney EB. The antidiuretic hormone and the factors which determine its release. *Proc R Soc Lond B Biol Sci.* 1947; **135**(878): 25-106.
3. Brooks FP, Pickford M. The effect of posterior pituitary hormones on the excretion of electrolytes, in dogs. *J Physiol.* 1958; **142**(3): 468-93.
4. Cross BA, Green JD. Activity of single neurones in the hypothalamus: effect of osmotic and other stimuli. *J Physiol.* 1959; **148**: 554-69.
5. Poulain DA, Wakerley JB, Dyball RE. Electrophysiological differentiation of oxytocin- and vasopressin-secreting neurones. *Proc R Soc Lond B Biol Sci.* 1977; **196**(1125): 367-84.
6. Leng G. Rat supraoptic neurones: the effects of locally applied hypertonic saline. *J Physiol.* 1980; **304**: 405-14.
7. Mason WT. Supraoptic neurones of rat hypothalamus are osmosensitive. *Nature.* 1980; **287**(5778): 154-7.
8. Gruber KA, Wilkin LD, Johnson AK. Neurohypophyseal hormone release and biosynthesis in rats with lesions of the anteroventral third ventricle (AV3V) region. *Brain Res.* 1986; **378**(1): 115-9.
9. McKinley MJ, Congiu M, Denton DA, Park RG, Penschow J, Simpson JB, Tarjan E, Weisinger RS, Wright RD. The anterior wall of the third cerebral ventricle and homeostatic responses to dehydration. *J Physiol (Paris).* 1984; **79**(6): 421-7.
10. Ramsay DJ, Thrasher TN, Keil LC. The organum vasculosum laminae terminalis: a critical area for osmoreception. *Prog Brain Res.* 1983; **60**: 1-8.
11. Johnson AK. The periventricular anteroventral third ventricle (AV3V): its relationship with the subfornical organ and neural systems involved in maintaining body fluid homeostasis. *Brain Res Bull.* 1985; **15**(6): 595-601.
12. Carithers J, Bealer SL, Brody MJ, Johnson AK. Fine structural evidence of degeneration in supraoptic nucleus and subfornical organ of rats with lesions in the anteroventral third ventricle. *Brain Res.* 1980; **201**(1): 1-12.
13. Mangiapane ML, Thrasher TN, Keil LC, Simpson JB, Ganong WF. Role for the subfornical organ in vasopressin release. *Brain Res Bull.* 1984; **13**(1): 43-7.
14. Prager-Khoutorsky M, Bourque CW. Anatomical organization of the rat organum vasculosum laminae terminalis. *Am J Physiol Regul Integr Comp Physiol.* 2015; **309**(4): R324-37.
15. McKinley MJ, Bicknell RJ, Hards D, McAllen RM, Vivas L, Weisinger RS, Oldfield BJ. Efferent neural pathways of the lamina terminalis subserving osmoregulation. *Prog Brain Res.* 1992; **91**: 395-402.
16. Rolls BJ. The effect of intravenous infusion of antidiuretic hormone on water intake in the rat. *J Physiol.* 1971; **219**(2): 331-9.
17. Park RG, Congiu M, Denton DA, McKinley MJ. Natriuresis induced by arginine vasopressin infusion in sheep. *Am J Physiol.* 1985; **249**(6 Pt 2): F799-805.
18. Lichardus B, McKinley MJ, Denton DA, McDougal JG, Weisinger RS, Okolicany J. On the brain involvement in saline loading natriuresis--an indirect evidence for the cerebral participation in the functional expression of the atrial natriuretic system. *Physiol Bohemoslov.* 1987; **36**(2): 175-8.
19. Balment RJ, Brimble MJ, Forsling ML. Release of oxytocin induced by salt loading and its influence on renal excretion in the male rat. *J Physiol.* 1980; **308**: 439-49.

20. Balment RJ, Brimble MJ, Forsling ML, Kelly LP, Musabayane CT. A synergistic effect of oxytocin and vasopressin on sodium excretion in the neurohypophysectomized rat. *J Physiol.* 1986; **381**:453-64.
21. Balment RJ, Brimble MJ, Forsling ML, Musabayane CT. The influence of neurohypophysial hormones on renal function in the acutely hypophysectomized rat. *J Physiol.* 1986; **381**:439-52.
22. Young TK, Van Dyke HB. Repletion of vasopressin and oxytocin in the posterior lobe of the pituitary gland of the rat. *J Endocrinol.* 1968; **40**(3): 337-42.
23. Williams TD, Abel DC, King CM, Jelley RY, Lightman SL. Vasopressin and oxytocin responses to acute and chronic osmotic stimuli in man. *J Endocrinol.* 1986; **108**(1): 163-8.
24. Conrad KP, Gellai M, North WG, Valtin H. Influence of oxytocin on renal hemodynamics and sodium excretion. *Ann N Y Acad Sci.* 1993; **689**:346-62.
25. Rasmussen MS, Simonsen JA, Sandgaard NC, Hoilund-Carlsen PF, Bie P. Effects of oxytocin in normal man during low and high sodium diets. *Acta Physiol Scand.* 2004; **181**(2): 247-57.
26. Bentley PJ. From a cat's blood pressure to a toad's bladder: pharmacology and the neurohypophysis--a review. *Mt Sinai J Med.* 1971; **38**(4): 344-62.
27. Lee J, deWardener HE. Neurosecretion and sodium excretion. *Kidney Int.* 1974; **6**(5): 323-30.
28. Brimble MJ, Dyball RE. Characterization of the responses of oxytocin- and vasopressin-secreting neurones in the supraoptic nucleus to osmotic stimulation. *J Physiol.* 1977; **271**(1): 253-71.
29. Brimble MJ, Dyball RE, Forsling ML. Oxytocin release following osmotic activation of oxytocin neurones in the paraventricular and supraoptic nuclei. *J Physiol.* 1978; **278**:69-78.
30. Conrad KP, Gellai M, North WG, Valtin H. Influence of oxytocin on renal hemodynamics and electrolyte and water excretion. *Am J Physiol.* 1986; **251**(2 Pt 2): F290-6.
31. Huang W, Lee SL, Arnason SS, Sjoquist M. Dehydration natriuresis in male rats is mediated by oxytocin. *Am J Physiol.* 1996; **270**(2 Pt 2): R427-33.
32. Windle RJ, Judah JM, Forsling ML. Effect of oxytocin receptor antagonists on the renal actions of oxytocin and vasopressin in the rat. *J Endocrinol.* 1997; **152**(2): 257-64.
33. Haanwinckel MA, Elias LK, Favaretto AL, Gutkowska J, McCann SM, Antunes-Rodrigues J. Oxytocin mediates atrial natriuretic peptide release and natriuresis after volume expansion in the rat. *Proc Natl Acad Sci U S A.* 1995; **92**(17): 7902-6.
34. McKinley MJ, Mathai ML, McAllen RM, McClear RC, Miselis RR, Pennington GL, Vivas L, Wade JD, Oldfield BJ. Vasopressin secretion: osmotic and hormonal regulation by the lamina terminalis. *J Neuroendocrinol.* 2004; **16**(4): 340-7.
35. Blackburn RE, Leng G, Russell JA. Control of magnocellular oxytocin neurones by the region anterior and ventral to the third ventricle (AV3V region) in rats. *J Endocrinol.* 1987; **114**(2): 253-61.
36. Russell JA, Blackburn RE, Leng G. Ablation of the region anterior and ventral to the third ventricle (AV3V region) does not impede parturition in rats. *J Endocrinol.* 1989; **121**(1): 109-15.
37. Meddle SL, Leng G, Selvarajah JR, Bicknell RJ, Russell JA. Direct pathways to the supraoptic nucleus from the brainstem and the main olfactory bulb are activated at parturition in the rat. *Neuroscience.* 2000; **101**(4): 1013-21.

38. Douglas A, Scullion S, Antonijevic I, Brown D, Russell J, Leng G. Uterine contractile activity stimulates supraoptic neurons in term pregnant rats via a noradrenergic pathway. *Endocrinology*. 2001; **142**(2): 633-44.
39. Russell JA, Leng G, Douglas AJ. The magnocellular oxytocin system, the fount of maternity: adaptations in pregnancy. *Front Neuroendocrinol*. 2003; **24**(1): 27-61.
40. Leng G, Mason WT, Dyer RG. The supraoptic nucleus as an osmoreceptor. *Neuroendocrinology*. 1982; **34**(1): 75-82.
41. Leng G, Brown CH, Bull PM, Brown D, Scullion S, Currie J, Blackburn-Munro RE, Feng J, Onaka T, Verbalis JG, Russell JA, Ludwig M. Responses of magnocellular neurons to osmotic stimulation involves coactivation of excitatory and inhibitory input: an experimental and theoretical analysis. *J Neurosci*. 2001; **21**(17): 6967-77.
42. MacGregor DJ, Leng G. Phasic firing in vasopressin cells: understanding its functional significance through computational models. *PLoS Comput Biol*. 2012; **8**(10): e1002740.
43. Douglass JK, Wilkens L, Pantazelou E, Moss F. Noise enhancement of information transfer in crayfish mechanoreceptors by stochastic resonance. *Nature*. 1993; **365**(6444): 337-40.
44. Oliet SH, Bourque CW. Mechanosensitive channels transduce osmosensitivity in supraoptic neurons. *Nature*. 1993; **364**(6435): 341-3.
45. Prager-Khoutorsky M, Khoutorsky A, Bourque CW. Unique interweaved microtubule scaffold mediates osmosensory transduction via physical interaction with TRPV1. *Neuron*. 2014; **83**(4): 866-78.
46. Bansal V, Fisher TE. Osmotic activation of a Ca(2+)-dependent phospholipase C pathway that regulates N TRPV1-mediated currents in rat supraoptic neurons. *Physiol Rep*. 2017; **5**(8).
47. Ciura S, Liedtke W, Bourque CW. Hypertonicity sensing in organum vasculosum lamina terminalis neurons: a mechanical process involving TRPV1 but not TRPV4. *J Neurosci*. 2011; **31**(41): 14669-76.
48. Tucker AB, Stocker SD. Hypernatremia-induced vasopressin secretion is not altered in TRPV1-/- rats. *Am J Physiol Regul Integr Comp Physiol*. 2016; **311**(3): R451-6.
49. Kinsman B, Cowles J, Lay J, Simmonds SS, Browning KN, Stocker SD. Osmoregulatory thirst in mice lacking the transient receptor potential vanilloid type 1 (TRPV1) and/or type 4 (TRPV4) receptor. *Am J Physiol Regul Integr Comp Physiol*. 2014; **307**(9): R1092-100.
50. Janas S, Seghers F, Schakman O, Alsady M, Deen P, Vriens J, Tissir F, Nilius B, Löffing J, Gailly P, Devuyst O. TRPV4 is associated with central rather than nephrogenic osmoregulation. *Pflugers Arch*. 2016; **468**(9): 1595-607.
51. O'Neil RG, Heller S. The mechanosensitive nature of TRPV channels. *Pflugers Arch*. 2005; **451**(1): 193-203.
52. Ciura S, Prager-Khoutorsky M, Thirouin ZS, Wyrosdic JC, Olson JE, Liedtke W, Bourque CW. Trpv4 Mediates Hypotonic Inhibition of Central Osmosensory Neurons via Taurine Gliotransmission. *Cell Rep*. 2018; **23**(8): 2245-53.
53. Nedungadi TP, Carreno FR, Walch JD, Bathina CS, Cunningham JT. Region-specific changes in transient receptor potential vanilloid channel expression in the vasopressin magnocellular system in hepatic cirrhosis-induced hyponatraemia. *J Neuroendocrinol*. 2012; **24**(4): 642-52.
54. Nedungadi TP, Dutta M, Bathina CS, Caterina MJ, Cunningham JT. Expression and distribution of TRPV2 in rat brain. *Exp Neurol*. 2012; **237**(1): 223-37.

55. Sladek CD, Johnson AK. Integration of thermal and osmotic regulation of water homeostasis: the role of TRPV channels. *Am J Physiol Regul Integr Comp Physiol*. 2013; **305**(7): R669-78.
56. Zaelzer C, Hua P, Prager-Khoutorsky M, Ciura S, Voisin DL, Liedtke W, Bourque CW. DeltaN-TRPV1: A Molecular Co-detector of Body Temperature and Osmotic Stress. *Cell Rep*. 2015; **13**(1): 23-30.
57. Tononi G, Sporns O, Edelman GM. Measures of degeneracy and redundancy in biological networks. *Proc Natl Acad Sci U S A*. 1999; **96**(6): 3257-62.
58. Tanaka M, Cummins TR, Ishikawa K, Black JA, Ibata Y, Waxman SG. Molecular and functional remodeling of electrogenic membrane of hypothalamic neurons in response to changes in their input. *Proc Natl Acad Sci U S A*. 1999; **96**(3): 1088-93.
59. Black JA, Vasylyev D, Dib-Hajj SD, Waxman SG. Nav1.9 expression in magnocellular neurosecretory cells of supraoptic nucleus. *Exp Neurol*. 2014; **253**: 174-9.
60. Sharma K, Haque M, Guidry R, Ueta Y, Teruyama R. Effect of dietary salt intake on epithelial Na(+) channels (ENaC) in vasopressin magnocellular neurosecretory neurons in the rat supraoptic nucleus. *J Physiol*. 2017; **595**(17): 5857-74.
61. Dine J, Ducourneau VR, Fenelon VS, Fossat P, Amadio A, Eder M, Israel JM, Oliet SH, Voisin DL. Extracellular signal-regulated kinase phosphorylation in forebrain neurones contributes to osmoregulatory mechanisms. *J Physiol*. 2014; **592**(7): 1637-54.
62. Prager-Khoutorsky M, Bourque CW. Mechanical basis of osmosensory transduction in magnocellular neurosecretory neurones of the rat supraoptic nucleus. *J Neuroendocrinol*. 2015; **27**(6): 507-15.
63. McKinley MJ, Allen AM, Burns P, Colvill LM, Oldfield BJ. Interaction of circulating hormones with the brain: the roles of the subfornical organ and the organum vasculosum of the lamina terminalis. *Clin Exp Pharmacol Physiol Suppl*. 1998; **25**: S61-7.
64. Sandgren JA, Linggongoro DW, Zhang SY, Sapouckey SA, Claflin KE, Pearson NA, Leidinger MR, Pierce GL, Santillan MK, Gibson-Corley KN, Sigmund CD, Grobe JL. Angiotensin AT1A receptors expressed in vasopressin-producing cells of the supraoptic nucleus contribute to osmotic control of vasopressin. *Am J Physiol Regul Integr Comp Physiol*. 2018; **314**(6): R770-R80.
65. Stachniak TJ, Trudel E, Bourque CW. Cell-specific retrograde signals mediate antiparallel effects of angiotensin II on osmoreceptor afferents to vasopressin and oxytocin neurons. *Cell Rep*. 2014; **8**(2): 355-62.
66. Chakfe Y, Bourque CW. Excitatory peptides and osmotic pressure modulate mechanosensitive cation channels in concert. *Nat Neurosci*. 2000; **3**(6): 572-9.
67. Ludwig M, Johnstone LE, Neumann I, Landgraf R, Russell JA. Direct hypertonic stimulation of the rat supraoptic nucleus increases c-fos expression in glial cells rather than magnocellular neurones. *Cell Tissue Res*. 1997; **287**(1): 79-90.
68. Deleuze C, Duvoid A, Hussy N. Properties and glial origin of osmotic-dependent release of taurine from the rat supraoptic nucleus. *J Physiol*. 1998; **507** (Pt 2): 463-71.
69. Choe KY, Olson JE, Bourque CW. Taurine release by astrocytes modulates osmosensitive glycine receptor tone and excitability in the adult supraoptic nucleus. *J Neurosci*. 2012; **32**(36): 12518-27.
70. Yang CR, Senatorov VV, Renaud LP. Organum vasculosum lamina terminalis-evoked postsynaptic responses in rat supraoptic neurones in vitro. *J Physiol*. 1994; **477**(Pt 1): 59-74.

71. Richard D, Bourque CW. Synaptic control of rat supraoptic neurones during osmotic stimulation of the organum vasculosum lamina terminalis in vitro. *J Physiol*. 1995; **489** (Pt 2)567-77.
72. Nissen R, Renaud LP. GABA receptor mediation of median preoptic nucleus-evoked inhibition of supraoptic neurosecretory neurones in rat. *J Physiol*. 1994; **479** (Pt 2)207-16.
73. Leng T, Leng G, MacGregor DJ. Spike patterning in oxytocin neurons: Capturing physiological behaviour with Hodgkin-Huxley and integrate-and-fire models. *PLoS One*. 2017; **12**(7): e0180368.
74. Sabatier N, Leng G. Bistability with hysteresis in the activity of vasopressin cells. *J Neuroendocrinol*. 2007; **19**(2): 95-101.
75. Ludwig M, Leng G. GABAergic projection from the arcuate nucleus to the supraoptic nucleus in the rat. *Neurosci Lett*. 2000; **281**(2-3): 195-7.
76. Haam J, Popescu IR, Morton LA, Halmos KC, Teruyama R, Ueta Y, Tasker JG. GABA is excitatory in adult vasopressinergic neuroendocrine cells. *J Neurosci*. 2012; **32**(2): 572-82.
77. Morton LA, Popescu IR, Haam J, Tasker JG. Short-term potentiation of GABAergic synaptic inputs to vasopressin and oxytocin neurones. *J Physiol*. 2014; **592**(19): 4221-33.
78. Choe KY, Trudel E, Bourque CW. Effects of Salt Loading on the Regulation of Rat Hypothalamic Magnocellular Neurosecretory Cells by Ionotropic GABA and Glycine Receptors. *J Neuroendocrinol*. 2016; **28**(4).
79. Korpak AK, Han SY, Schwenke DO, Brown CH. A switch from GABA inhibition to excitation of vasopressin neurons exacerbates the development angiotensin II-dependent hypertension. *J Neuroendocrinol*. 2017.
80. Kim YB, Kim WB, Jung WW, Jin X, Kim YS, Kim B, Han HC, Block GD, Colwell CS, Kim YI. Excitatory GABAergic Action and Increased Vasopressin Synthesis in Hypothalamic Magnocellular Neurosecretory Cells Underlie the High Plasma Level of Vasopressin in Diabetic Rats. *Diabetes*. 2018; **67**(3): 486-95.
81. Lee SW, Kim YB, Kim JS, Kim WB, Kim YS, Han HC, Colwell CS, Cho YW, In Kim Y. GABAergic inhibition is weakened or converted into excitation in the oxytocin and vasopressin neurons of the lactating rat. *Mol Brain*. 2015; **8**34.
82. Srisawat R, Ludwig M, Bull PM, Douglas AJ, Russell JA, Leng G. Nitric oxide and the oxytocin system in pregnancy. *J Neurosci*. 2000; **20**(17): 6721-7.
83. Sabatier N, Leng G. Presynaptic actions of endocannabinoids mediate alpha-MSH-induced inhibition of oxytocin cells. *Am J Physiol Regul Integr Comp Physiol*. 2006; **290**(3): R577-84.
84. Srisawat R, Bishop VR, Bull PM, Douglas AJ, Russell JA, Ludwig M, Leng G. Regulation of neuronal nitric oxide synthase mRNA expression in the rat magnocellular neurosecretory system. *Neurosci Lett*. 2004; **369**(3): 191-6.
85. Armstrong W Eea. Electrophysiological Properties of Identified Oxytocin and Vasopressin Neurons. *Journal of Neuroendocrinology*. 2019; **in press**.
86. da Silva MP, Merino RM, Mecawi AS, Moraes DJ, Varanda WA. In vitro differentiation between oxytocin- and vasopressin-secreting magnocellular neurons requires more than one experimental criterion. *Mol Cell Endocrinol*. 2015; **400**102-11.
87. Bicknell RJ. Optimizing release from peptide hormone secretory nerve terminals. *J Exp Biol*. 1988; **139**51-65.
88. Verbalis JG, Mangione MP, Stricker EM. Oxytocin produces natriuresis in rats at physiological plasma concentrations. *Endocrinology*. 1991; **128**(3): 1317-22.

89. Sjoquist M, Huang W, Jacobsson E, Skott O, Stricker EM, Sved AF. Sodium excretion and renin secretion after continuous versus pulsatile infusion of oxytocin in rats. *Endocrinology*. 1999; **140**(6): 2814-8.
90. Stricker EM, Thiels E, Verbalis JG. Sodium appetite in rats after prolonged dietary sodium deprivation: a sexually dimorphic phenomenon. *Am J Physiol*. 1991; **260**(6 Pt 2): R1082-8.
91. Richter CP. Increased salt appetite in adrenalectomized rats. *American Journal of Physiology*. 1936; **115**(1): 155-61.
92. Bykowski MR, Smith JC, Stricker EM. Regulation of NaCl solution intake and gastric emptying in adrenalectomized rats. *Physiol Behav*. 2007; **92**(5): 781-9.
93. Gasparini S, Melo MR, Leite GF, Nascimento PA, Andrade-Franze GMF, Menani JV, Colombari E. Rapid stimulation of sodium intake combining aldosterone into the 4th ventricle and the blockade of the lateral parabrachial nucleus. *Neuroscience*. 2017; **346**:94-101.
94. Joels M, de Kloet ER. 30 YEARS OF THE MINERALOCORTICOID RECEPTOR: The brain mineralocorticoid receptor: a saga in three episodes. *J Endocrinol*. 2017; **234**(1): T49-T66.
95. Stricker EM, Jalowiec JE. Restoration of Intravascular Fluid Volume Following Acute Hypovolemia in Rats. *American Journal of Physiology*. 1970; **218**(1): 191-&.
96. Smith CA, Curtis KS, Smith JC, Stricker EM. Presystemic influences on thirst, salt appetite, and vasopressin secretion in the hypovolemic rat. *Am J Physiol Regul Integr Comp Physiol*. 2007; **292**(5): R2089-99.
97. Stricker EM, Hoffmann ML. Presystemic signals in the control of thirst, salt appetite, and vasopressin secretion. *Physiol Behav*. 2007; **91**(4): 404-12.
98. McKinley MJ, Walker LL, Alexiou T, Allen AM, Campbell DJ, Di Nicolantonio R, Oldfield BJ, Denton DA. Osmoregulatory fluid intake but not hypovolemic thirst is intact in mice lacking angiotensin. *Am J Physiol Regul Integr Comp Physiol*. 2008; **294**(5): R1533-43.
99. Matsuda T, Hiyama TY, Niimura F, Matsusaka T, Fukamizu A, Kobayashi K, Kobayashi K, Noda M. Distinct neural mechanisms for the control of thirst and salt appetite in the subfornical organ. *Nat Neurosci*. 2017; **20**(2): 230-41.
100. Nation HL, Nicoleau M, Kinsman BJ, Browning KN, Stocker SD. DREADD-induced activation of subfornical organ neurons stimulates thirst and salt appetite. *J Neurophysiol*. 2016; **115**(6): 3123-9.
101. Hurley SW, Zhang Z, Beltz TG, Xue B, Johnson AK. Sensitization of sodium appetite: evidence for sustained molecular changes in the lamina terminalis. *Am J Physiol Regul Integr Comp Physiol*. 2014; **307**(12): R1405-12.
102. Roncari CF, Barbosa RM, Vendramini RC, De Luca LA, Jr., Menani JV, Colombari E, Colombari DSA. Enhanced angiotensin II induced sodium appetite in renovascular hypertensive rats. *Peptides*. 2018; **101**:82-8.
103. Stricker EM, Verbalis JG. Inhibition of salt appetite in rats by central oxytocin. *Am J Physiol Regul Integr Comp Physiol*. 2004; **287**(2): R487; author reply R-8.
104. Grafe LA, Takacs AE, Yee DK, Flanagan-Cato LM. The role of the hypothalamic paraventricular nucleus and the organum vasculosum lateral terminalis in the control of sodium appetite in male rats. *J Neurosci*. 2014; **34**(28): 9249-60.
105. Blackburn RE, Stricker EM, Verbalis JG. Acute effects of ethanol on ingestive behavior in rats. *Alcohol Clin Exp Res*. 1994; **18**(4): 924-30.

106. Blackburn RE, Samson WK, Fulton RJ, Stricker EM, Verbalis JG. Central oxytocin inhibition of salt appetite in rats: evidence for differential sensing of plasma sodium and osmolality. *Proc Natl Acad Sci U S A*. 1993; **90**(21): 10380-4.
107. Blackburn RE, Samson WK, Fulton RJ, Stricker EM, Verbalis JG. Central oxytocin and ANP receptors mediate osmotic inhibition of salt appetite in rats. *Am J Physiol*. 1995; **269**(2 Pt 2): R245-51.
108. Fitts DA, Thornton SN, Ruhf AA, Zierath DK, Johnson AK, Thunhorst RL. Effects of central oxytocin receptor blockade on water and saline intake, mean arterial pressure, and c-Fos expression in rats. *Am J Physiol Regul Integr Comp Physiol*. 2003; **285**(6): R1331-9.
109. White MM, Samson WK. A possible relationship between brain-derived adrenomedullin and oxytocin in the regulation of sodium balance. *J Endocrinol*. 2009; **203**(2): 253-62.
110. Inyushkin AN, Orlans HO, Dyball RE. Secretory cells of the supraoptic nucleus have central as well as neurohypophysial projections. *J Anat*. 2009; **215**(4): 425-34.
111. Ryan PJ, Ross SI, Campos CA, Derkach VA, Palmiter RD. Oxytocin-receptor-expressing neurons in the parabrachial nucleus regulate fluid intake. *Nat Neurosci*. 2017; **20**(12): 1722-33.
112. Godino A, Margatho LO, Caeiro XE, Antunes-Rodrigues J, Vivas L. Activation of lateral parabrachial afferent pathways and endocrine responses during sodium appetite regulation. *Exp Neurol*. 2010; **221**(2): 275-84.
113. Johnstone LE, Fong TM, Leng G. Neuronal activation in the hypothalamus and brainstem during feeding in rats. *Cell Metab*. 2006; **4**(4): 313-21.
114. Mandelblat-Cerf Y, Kim A, Burgess CR, Subramanian S, Tannous BA, Lowell BB, Andermann ML. Bidirectional Anticipation of Future Osmotic Challenges by Vasopressin Neurons. *Neuron*. 2017; **93**(1): 57-65.
115. Leng G, Sabatier N. Oxytocin - The Sweet Hormone? *Trends Endocrinol Metab*. 2017; **28**(5): 365-76.
116. Song Z, Levin BE, Stevens W, Sladek CD. Supraoptic oxytocin and vasopressin neurons function as glucose and metabolic sensors. *Am J Physiol Regul Integr Comp Physiol*. 2014; **306**(7): R447-56.
117. Sabatier N, Caqueneau C, Dayanithi G, Bull P, Douglas AJ, Guan XM, Jiang M, Van der Ploeg L, Leng G. Alpha-melanocyte-stimulating hormone stimulates oxytocin release from the dendrites of hypothalamic neurons while inhibiting oxytocin release from their terminals in the neurohypophysis. *J Neurosci*. 2003; **23**(32): 10351-8.
118. Velmurugan S, Brunton PJ, Leng G, Russell JA. Circulating secretin activates supraoptic nucleus oxytocin and vasopressin neurons via noradrenergic pathways in the rat. *Endocrinology*. 2010; **151**(6): 2681-8.
119. Velmurugan S, Russell JA, Leng G. Systemic leptin increases the electrical activity of supraoptic nucleus oxytocin neurones in virgin and late pregnant rats. *J Neuroendocrinol*. 2013; **25**(4): 383-90.
120. Onaka T, Luckman SM, Antonijevic I, Palmer JR, Leng G. Involvement of the noradrenergic afferents from the nucleus tractus solitarius to the supraoptic nucleus in oxytocin release after peripheral cholecystokinin octapeptide in the rat. *Neuroscience*. 1995; **66**(2): 403-12.
121. Ong ZY, Bongiorno DM, Hernando MA, Grill HJ. Effects of Endogenous Oxytocin Receptor Signaling in Nucleus Tractus Solitarius on Satiation-Mediated Feeding and Thermogenic Control in Male Rats. *Endocrinology*. 2017; **158**(9): 2826-36.

122. Gartner SN, Aidney F, Klockars A, Prosser C, Carpenter EA, Isgrove K, Levine AS, Olszewski PK. Intra gastric preloads of l-tryptophan reduce ingestive behavior via oxytocinergic neural mechanisms in male mice. *Appetite*. 2018; **125**:278-86.
123. Hume C, Sabatier N, Menzies J. High-Sugar, but Not High-Fat, Food Activates Supraoptic Nucleus Neurons in the Male Rat. *Endocrinology*. 2017; **158**(7): 2200-11.
124. Kohno D, Nakata M, Maejima Y, Shimizu H, Sedbazar U, Yoshida N, Dezaki K, Onaka T, Mori M, Yada T. Nesfatin-1 neurons in paraventricular and supraoptic nuclei of the rat hypothalamus coexpress oxytocin and vasopressin and are activated by refeeding. *Endocrinology*. 2008; **149**(3): 1295-301.
125. Saito R, So M, Motojima Y, Matsuura T, Yoshimura M, Hashimoto H, Yamamoto Y, Kusuhara K, Ueta Y. Activation of Nesfatin-1-Containing Neurons in the Hypothalamus and Brainstem by Peripheral Administration of Anorectic Hormones and Suppression of Feeding via Central Nesfatin-1 in Rats. *J Neuroendocrinol*. 2016; **28**(9).
126. Palkovits M, Kiss JZ, Beinfeld MC, Brownstein MJ. Cholecystokinin in the hypothalamo-hypophyseal system. *Brain Res*. 1984; **299**(1): 186-9.
127. Shen J, Larm JA, Gundlach AL. Galanin-like peptide mRNA in neural lobe of rat pituitary. Increased expression after osmotic stimulation suggests a role for galanin-like peptide in neuron-glial interactions and/or neurosecretion. *Neuroendocrinology*. 2001; **73**(1): 2-11.
128. Gillard ER, Leon-Olea M, Mucio-Ramirez S, Coburn CG, Sanchez-Islas E, de Leon A, Mussenden H, Bauce LG, Pittman QJ, Curras-Collazo MC. A novel role for endogenous pituitary adenylate cyclase activating polypeptide in the magnocellular neuroendocrine system. *Endocrinology*. 2006; **147**(2): 791-803.
129. Sakai RR, McEwen BS, Fluharty SJ, Ma LY. The amygdala: site of genomic and nongenomic arousal of aldosterone-induced sodium intake. *Kidney Int*. 2000; **57**(4): 1337-45.
130. Klockars OA, Klockars A, Levine AS, Olszewski PK. Oxytocin administration in the basolateral and central nuclei of amygdala moderately suppresses food intake. *Neuroreport*. 2018; **29**(6): 504-10.
131. Knobloch HS, Charlet A, Hoffmann LC, Eliava M, Khrulev S, Cetin AH, Osten P, Schwarz MK, Seeburg PH, Stoop R, Grinevich V. Evoked axonal oxytocin release in the central amygdala attenuates fear response. *Neuron*. 2012; **73**(3): 553-66.
132. Ramsay DS, Woods SC. Clarifying the roles of homeostasis and allostasis in physiological regulation. *Psychol Rev*. 2014; **121**(2): 225-47.
133. Stricker EM, Callahan JB, Huang W, Sved AF. Early osmoregulatory stimulation of neurohypophyseal hormone secretion and thirst after gastric NaCl loads. *Am J Physiol Regul Integr Comp Physiol*. 2002; **282**(6): R1710-7.
134. Stricker EM, Huang W, Sved AF. Early osmoregulatory signals in the control of water intake and neurohypophyseal hormone secretion. *Physiol Behav*. 2002; **76**(3): 415-21.
135. Gizowski C, Bourque CW. The neural basis of homeostatic and anticipatory thirst. *Nat Rev Nephrol*. 2018; **14**(1): 11-25.
136. Trudel E, Bourque CW. Circadian modulation of osmoregulated firing in rat supraoptic nucleus neurones. *J Neuroendocrinol*. 2012; **24**(4): 577-86.
137. Qiu J, Hindmarch CC, Yao ST, Tasker JG, Murphy D. Transcriptomic analysis of the osmotic and reproductive remodeling of the female rat supraoptic nucleus. *Endocrinology*. 2011; **152**(9): 3483-91.

138. Johnson KR, Hindmarch CC, Salinas YD, Shi Y, Greenwood M, Hoe SZ, Murphy D, Gainer H. A RNA-Seq Analysis of the Rat Supraoptic Nucleus Transcriptome: Effects of Salt Loading on Gene Expression. *PLoS One*. 2015; **10**(4): e0124523.
139. Greenwood MP, Mecawi AS, Hoe SZ, Mustafa MR, Johnson KR, Al-Mahmoud GA, Elias LL, Paton JF, Antunes-Rodrigues J, Gainer H, Murphy D, Hindmarch CC. A comparison of physiological and transcriptome responses to water deprivation and salt loading in the rat supraoptic nucleus. *Am J Physiol Regul Integr Comp Physiol*. 2015; **308**(7): R559-68.
140. Tasker JG, Oliet SH, Bains JS, Brown CH, Stern JE. Glial regulation of neuronal function: from synapse to systems physiology. *J Neuroendocrinol*. 2012; **24**(4): 566-76.
141. Reis WL, Biancardi VC, Son S, Antunes-Rodrigues J, Stern JE. Carbon monoxide and nitric oxide interactions in magnocellular neurosecretory neurones during water deprivation. *J Neuroendocrinol*. 2015; **27**(2): 111-22.
142. Ladyman SR, Khant Aung Z, Grattan DR. Impact of Pregnancy and Lactation on the Long-Term Regulation of Energy Balance in Female Mice. *Endocrinology*. 2018; **159**(6): 2324-36.
143. Augustine RA, Ladyman SR, Bouwer GT, Alyousif Y, Sapsford TJ, Scott V, Kokay IC, Grattan DR, Brown CH. Prolactin regulation of oxytocin neurone activity in pregnancy and lactation. *J Physiol*. 2017; **595**(11): 3591-605.
144. Augustine RA, Bouwer GT, Seymour AJ, Grattan DR, Brown CH. Reproductive Regulation of Gene Expression in the Hypothalamic Supraoptic and Paraventricular Nuclei. *J Neuroendocrinol*. 2016; **28**(4).
145. Augustine RA, Seymour AJ, Campbell RE, Grattan DR, Brown CH. Integrative neuro-humoral regulation of oxytocin neuron activity in pregnancy and lactation. *J Neuroendocrinol*. 2018.
146. Seymour AJ, Scott V, Augustine RA, Bouwer GT, Campbell RE, Brown CH. Development of an excitatory kisspeptin projection to the oxytocin system in late pregnancy. *J Physiol*. 2017; **595**(3): 825-38.
147. Sunn N, Egli M, Burazin TC, Burns P, Colvill L, Davern P, Denton DA, Oldfield BJ, Weisinger RS, Rauch M, Schmid HA, McKinley MJ. Circulating relaxin acts on subfornical organ neurons to stimulate water drinking in the rat. *Proc Natl Acad Sci U S A*. 2002; **99**(3): 1701-6.
148. Atherton JC, Dark JM, Garland HO, Morgan MR, Pidgeon J, Soni S. Changes in water and electrolyte balance, plasma volume and composition during pregnancy in the rat. *J Physiol*. 1982; **330**: 81-93.
149. Brunton PJ, Arunachalam S, Russel JA. Control of neurohypophysial hormone secretion, blood osmolality and volume in pregnancy. *J Physiol Pharmacol*. 2008; **59 Suppl**: 827-45.
150. Way SA, Leng G. Relaxin increases the firing rate of supraoptic neurones and increases oxytocin secretion in the rat. *J Endocrinol*. 1992; **132**(1): 149-58.
151. Brown CH, Bains JS, Ludwig M, Stern JE. Physiological regulation of magnocellular neurosecretory cell activity: integration of intrinsic, local and afferent mechanisms. *J Neuroendocrinol*. 2013; **25**(8): 678-710.
152. Douglas AJ, Dye S, Leng G, Russell JA, Bicknell RJ. Endogenous opioid regulation of oxytocin secretion through pregnancy in the rat. *J Neuroendocrinol*. 1993; **5**(3): 307-14.
153. Leng G, Dye S, Bicknell RJ. Kappa-opioid restraint of oxytocin secretion: plasticity through pregnancy. *Neuroendocrinology*. 1997; **66**(6): 378-83.

154. Russell JA, Douglas AJ, Bull PM, Pumford KM, Bicknell RJ, Leng G. Pregnancy and opioid interactions with the anterior perithird ventricular input to magnocellular oxytocin neurones. *Prog Brain Res*. 1992; **91**:141-53.
155. Douglas AJ, Neumann I, Meeren HK, Leng G, Johnstone LE, Munro G, Russell JA. Central endogenous opioid inhibition of supraoptic oxytocin neurons in pregnant rats. *J Neurosci*. 1995; **15**(7 Pt 1): 5049-57.
156. Douglas AJ, Bicknell RJ, Leng G, Russell JA, Meddle SL. Beta-endorphin cells in the arcuate nucleus: projections to the supraoptic nucleus and changes in expression during pregnancy and parturition. *J Neuroendocrinol*. 2002; **14**(10): 768-77.
157. Ladyman SR, Augustine RA, Grattan DR. Hormone interactions regulating energy balance during pregnancy. *J Neuroendocrinol*. 2010; **22**(7): 805-17.
158. Douglas AJ, Johnstone LE, Leng G. Neuroendocrine mechanisms of change in food intake during pregnancy: a potential role for brain oxytocin. *Physiol Behav*. 2007; **91**(4): 352-65.
159. Onaka T, Luckman SM, Guevara-Guzman R, Ueta Y, Kendrick K, Leng G. Presynaptic actions of morphine: blockade of cholecystokinin-induced noradrenaline release in the rat supraoptic nucleus. *J Physiol*. 1995; **482** (Pt 1):69-79.
160. Brunton PJ, Meddle SL, Ma S, Ochedalski T, Douglas AJ, Russell JA. Endogenous opioids and attenuated hypothalamic-pituitary-adrenal axis responses to immune challenge in pregnant rats. *J Neurosci*. 2005; **25**(21): 5117-26.
161. Shibuki K, Leng G, Way S. Effects of naloxone and of intraperitoneal hypertonic saline upon oxytocin release and upon supraoptic neuronal activity. *Neurosci Lett*. 1988; **88**(1): 75-80.
162. Francis K, Meddle SL, Bishop VR, Russell JA. Progesterone receptor expression in the pregnant and parturient rat hypothalamus and brainstem. *Brain Res*. 2002; **927**(1): 18-26.
163. Brussaard AB, Herbison AE. Long-term plasticity of postsynaptic GABAA-receptor function in the adult brain: insights from the oxytocin neurone. *Trends Neurosci*. 2000; **23**(5): 190-5.
164. Koksma JJ, van Kesteren RE, Rosahl TW, Zwart R, Smit AB, Luddens H, Brussaard AB. Oxytocin regulates neurosteroid modulation of GABA(A) receptors in supraoptic nucleus around parturition. *J Neurosci*. 2003; **23**(3): 788-97.
165. Brunton PJ, McKay AJ, Ochedalski T, Piastowska A, Rebas E, Lachowicz A, Russell JA. Central opioid inhibition of neuroendocrine stress responses in pregnancy in the rat is induced by the neurosteroid allopregnanolone. *J Neurosci*. 2009; **29**(20): 6449-60.
166. Brunton PJ, Sabatier N, Leng G, Russell JA. Suppressed oxytocin neuron responses to immune challenge in late pregnant rats: a role for endogenous opioids. *Eur J Neurosci*. 2006; **23**(5): 1241-7.
167. Brunton PJ, Bales J, Russell JA. Allopregnanolone and Induction of Endogenous Opioid Inhibition of Oxytocin Responses to Immune Stress in Pregnant Rats. *Journal of Neuroendocrinology*. 2012; **24**(4): 690-700.
168. Eliava M, Melchior M, Knobloch-Bollmann HS, Wahis J, da Silva Gouveia M, Tang Y, Ciobanu AC, Triana Del Rio R, Roth LC, Althammer F, Chavant V, Goumon Y, Gruber T, Petit-Demouliere N, Busnelli M, Chini B, Tan LL, Mitre M, Froemke RC, Chao MV, Giese G, Sprengel R, Kuner R, Poisbeau P, Seeburg PH, Stoop R, Charlet A, Grinevich V. A New Population of Parvocellular Oxytocin Neurons Controlling Magnocellular Neuron Activity and Inflammatory Pain Processing. *Neuron*. 2016; **89**(6): 1291-304.

169. Juif PE, Breton JD, Rajalu M, Charlet A, Goumon Y, Poisbeau P. Long-lasting spinal oxytocin analgesia is ensured by the stimulation of allopregnanolone synthesis which potentiates GABA(A) receptor-mediated synaptic inhibition. *J Neurosci*. 2013; **33**(42): 16617-26.
170. Liutkeviciute Z, Koehbach J, Eder T, Gil-Mansilla E, Gruber CW. Global map of oxytocin/vasopressin-like neuropeptide signalling in insects. *Sci Rep*. 2016; **6**: 39177.
171. Minakata H. Oxytocin/vasopressin and gonadotropin-releasing hormone from cephalopods to vertebrates. *Ann N Y Acad Sci*. 2010; **1200**: 33-42.
172. Lamm MS, Liu H, Gemmell NJ, Godwin JR. The Need for Speed: Neuroendocrine Regulation of Socially-controlled Sex Change. *Integr Comp Biol*. 2015; **55**(2): 307-22.
173. Yamashita J, Kawabata Y, Okubo K. Expression of isotocin is male-specifically up-regulated by gonadal androgen in the medaka brain. *J Neuroendocrinol*. 2017; **29**(12).
174. Gesto M, Soengas JL, Rodriguez-Illamola A, Miguez JM. Arginine vasotocin treatment induces a stress response and exerts a potent anorexigenic effect in rainbow trout, *Oncorhynchus mykiss*. *J Neuroendocrinol*. 2014; **26**(2): 89-99.
175. Mennigen JA, Volkoff H, Chang JP, Trudeau VL. The nonapeptide isotocin in goldfish: Evidence for serotonergic regulation and functional roles in the control of food intake and pituitary hormone release. *Gen Comp Endocrinol*. 2017; **254**: 38-49.
176. Lagman D, Ocampo Daza D, Widmark J, Abalo XM, Sundstrom G, Larhammar D. The vertebrate ancestral repertoire of visual opsins, transducin alpha subunits and oxytocin/vasopressin receptors was established by duplication of their shared genomic region in the two rounds of early vertebrate genome duplications. *BMC Evol Biol*. 2013; **13**: 238.
177. Mayasich SA, Clarke BL. The emergence of the vasopressin and oxytocin hormone receptor gene family lineage: Clues from the characterization of vasotocin receptors in the sea lamprey (*Petromyzon marinus*). *Gen Comp Endocrinol*. 2016; **226**: 88-101.
178. Tessmar-Raible K, Raible F, Christodoulou F, Guy K, Rembold M, Hausen H, Arendt D. Conserved sensory-neurosecretory cell types in annelid and fish forebrain: insights into hypothalamus evolution. *Cell*. 2007; **129**(7): 1389-400.
179. Murphy D, Si-Hoe SL, Brenner S, Venkatesh B. Something fishy in the rat brain: molecular genetics of the hypothalamo-neurohypophysial system. *Bioessays*. 1998; **20**(9): 741-9.
180. Beets I, Janssen T, Meelkop E, Temmerman L, Suetens N, Rademakers S, Jansen G, Schoofs L. Vasopressin/Oxytocin-Related Signaling Regulates Gustatory Associative Learning in *C. elegans*. *Science*. 2012; **338**(6106): 543-5.
181. Hrabovszky E, Csapo AK, Kallo I, Wilhelm T, Turi GF, Liposits Z. Localization and osmotic regulation of vesicular glutamate transporter-2 in magnocellular neurons of the rat hypothalamus. *Neurochem Int*. 2006; **48**(8): 753-61.
182. Althammer F, Grinevich V. Diversity of oxytocin neurons: beyond magno- and parvocellular cell types? *J Neuroendocrinol*. 2017.
183. Chini B, Verhage M, Grinevich V. The Action Radius of Oxytocin Release in the Mammalian CNS: From Single Vesicles to Behavior. *Trends Pharmacol Sci*. 2017; **38**(11): 982-91.
184. Herisson FM, Waas JR, Fredriksson R, Schioth HB, Levine AS, Olszewski PK. Oxytocin Acting in the Nucleus Accumbens Core Decreases Food Intake. *J Neuroendocrinol*. 2016; **28**(4).
185. Dumais KM, Alonso AG, Bredewold R, Veenema AH. Role of the oxytocin system in amygdala subregions in the regulation of social interest in male and female rats. *Neuroscience*. 2016; **330**: 138-49.

186. Dumais KM, Alonso AG, Immormino MA, Bredewold R, Veenema AH. Involvement of the oxytocin system in the bed nucleus of the stria terminalis in the sex-specific regulation of social recognition. *Psychoneuroendocrinology*. 2016; **64**:79-88.
187. Moaddab M, Dabrowska J. Oxytocin receptor neurotransmission in the dorsolateral bed nucleus of the stria terminalis facilitates the acquisition of cued fear in the fear-potentiated startle paradigm in rats. *Neuropharmacology*. 2017; **121**:130-9.
188. Raam T, McAvoy KM, Besnard A, Veenema AH, Sahay A. Hippocampal oxytocin receptors are necessary for discrimination of social stimuli. *Nat Commun*. 2017; **8**(1): 2001.
189. Ludwig M, Leng G. Dendritic peptide release and peptide-dependent behaviours. *Nat Rev Neurosci*. 2006; **7**(2): 126-36.
190. Rossoni E, Feng J, Tirozzi B, Brown D, Leng G, Moos F. Emergent synchronous bursting of oxytocin neuronal network. *PLoS Comput Biol*. 2008; **4**(7): e1000123.
191. Leng G, Ludwig M. Neurotransmitters and peptides: whispered secrets and public announcements. *J Physiol*. 2008; **586**(23): 5625-32.
192. Yoshimura M, Nishimura K, Nishimura H, Sonoda S, Ueno H, Motojima Y, Saito R, Maruyama T, Nonaka Y, Ueta Y. Activation of endogenous arginine vasopressin neurons inhibit food intake: by using a novel transgenic rat line with DREADDs system. *Sci Rep*. 2017; **7**(1): 15728.
193. Pei H, Sutton AK, Burnett KH, Fuller PM, Olson DP. AVP neurons in the paraventricular nucleus of the hypothalamus regulate feeding. *Mol Metab*. 2014; **3**(2): 209-15.
194. Leng G, Pineda R, Sabatier N, Ludwig M. 60 YEARS OF NEUROENDOCRINOLOGY: The posterior pituitary, from Geoffrey Harris to our present understanding. *J Endocrinol*. 2015; **226**(2): T173-85.
195. Tobin VA, Hashimoto H, Wacker DW, Takayanagi Y, Langnaese K, Caquineau C, Noack J, Landgraf R, Onaka T, Leng G, Meddle SL, Engelmann M, Ludwig M. An intrinsic vasopressin system in the olfactory bulb is involved in social recognition. *Nature*. 2010; **464**(7287): 413-7.
196. Tsuji T, Allchorne AJ, Zhang M, Tsuji C, Tobin VA, Pineda R, Raftogianni A, Stern JE, Grinevich V, Leng G, Ludwig M. Vasopressin casts light on the suprachiasmatic nucleus. *J Physiol*. 2017; **595**(11): 3497-514.

Figure Legends

Figure 1. The osmoresponsiveness of magnocellular neurones.

A. Response of a vasopressin cell of the rat supraoptic nucleus to systemic osmotic stimulation in a urethane-anaesthetised rat. The graph plots the firing rate in 1-min bins during continuous i.v. infusion of 1 M NaCl at 26 μ l/min for more than 2 h. The protocol was similar to that in Leng et al. (41), but a lower concentration of NaCl was used over a longer time. The data are from unpublished experiments with Nancy Sabatier used to support the development of a computational model (42). The extracts of spike activity in B-D are from at the beginning of infusion (B); after 4000 s (C); and after 8000 s (D). They show the evolution of intense phasic activity in C, and then of continuous fast spiking activity in D. The minute-by-minute variability

of firing rate reflects the intermittent phasic firing pattern, but note the linear rise in mean rate over the duration of infusion.

E. The osmosensitiveness of magnocellular neurons involves an increase in afferent input and a direct osmosensitive mechanism. Here in a simple simulation, a fluctuating membrane potential around a mean level 9 mV below the spike threshold (indicated by the red lines) is mimicked by a randomly generated sequence of simplified EPSPs and IPSPs, each with an amplitude of 3 mV and a half-life of 5 ms, arriving at an equal mean rate of 100 Hz. None of the fluctuations cross the spike threshold. If the spike threshold is reduced by 2 mV (green line) mimicking the effects of a 2 mV direct depolarization, just one of the fluctuations crosses the threshold (green asterisk). Thus at this level of synaptic input, a small direct depolarization has little effect on spiking activity.

F, In this case, the mean input rate for both EPSPs and IPSPs is 200 Hz. Now, the fluctuations exceed the spike threshold on three occasions (red asterisks), and a depolarization of 2 mV results in an additional 11 threshold crossings. Thus in the presence of sufficient synaptic input, small levels of direct depolarisation can have a large effect on the spiking activity of magnocellular neurones.

Figure 2. GABA inputs to vasopressin cells.

In these experiments, single vasopressin cells were recorded from the supraoptic nucleus of a urethane-anaesthetised rat, and single electrical stimuli were applied to the OVLT every 5 s.

A. The experimental set-up. The supraoptic nucleus was exposed by ventral surgery, a stimulating electrode was placed on the neural stalk to allow supraoptic neurons to be antidromically identified, a microdialysis loop was placed on the ventral surface of the nucleus to allow direct application of the GABA antagonist bicuculline, and a stimulating electrode was placed on the OVLT.

B shows interspike interval (ISI) distributions compiled over 1200 s of activity before (orange) and after (blue) bicuculline, showing the resultant increase in mean activity.

C shows mean spike activity before (above) and below (during) bicuculline.

D shows the response to OVLT stimulation before (orange) and after (blue) bicuculline as post-stimulus time histograms (in 1-ms bins) constructed over the periods shown in D. In this cell,

OVLT stimulation produced a marked inhibition; bicuculline blocked this inhibition and unmasked an excitatory response. See (74) for details.

E shows the difference between the two histograms in D, showing the inferred inhibitory effects of OVLT stimulation by subtracting the excitatory effects (blue in D) from the mixed effects (orange in D), indicating that in this cell, the inhibitory effects of OVLT stimulation have a latency of onset of about 20-30 ms.

Figure 3. The main regulators of osmoresponsiveness in oxytocin cells.

In the rat, extracellular osmotic pressure and increased $[Na^+]$ are sensed both by magnocellular neurons and by cells in the subfornical organ (SFO) and OVLT. Cells in the SFO and OVLT are also responsive to changes in the circulating levels of a number of blood-borne hormones that have important effects on fluid and electrolyte homeostasis. Cells in the OVLT and SFO project directly to the magnocellular neurons, and this direct projection involves the excitatory transmitter glutamate and various peptides. They also project indirectly via the nucleus medianus, and this projection involves the inhibitory transmitter GABA. Astrocytes in the supraoptic nucleus release taurine in response to hypotonic stimulation, and this inhibits via action at glycine receptors on the supraoptic neurons. The GABA-ergic inputs are amplified by nitric oxide, produced by oxytocin cells in an activity-dependent manner (82), and the glutamatergic inputs are moderated by endocannabinoids, also produced by oxytocin cells in an activity-dependent manner (83). Oxytocin release is autoregulated by dynorphin at the level of the nerve terminals in the pituitary. In pregnancy, allopregnanolone enhances the inhibitory effects of GABA, and there is upregulation of both dynorphin expression and down regulation of nitric oxide synthase activity (84).

Figure 4. Oxytocin cell responses in pregnancy

A. In pregnancy, endogenous opioids suppress oxytocin secretion. In these experiments we blocked all opioid actions by pretreating the rats with naloxone to assess opioid-independent changes in the responsiveness of the oxytocin system. Responses were measured in virgin (black bars) and late pregnant rats (day 21, blue bars). The bars show increases in plasma oxytocin concentration (S.E.M.) above basal levels in response to different stimuli. Changes were measured in anaesthetized rats after electrical stimulation of the AV3V region; hypertonic saline,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

CCK; and in conscious rats following IL-1 β . Both AV3V stimulation and hypertonic saline stimulated oxytocin secretion less effectively in late pregnant rats. Conversely, oxytocin secretory responses to CCK and IL-1 β are greater in late pregnant rats. See (149) for details.

B. Responses of oxytocin cells to CCK (firing rate in 1-min bins above mean basal level) were measured in the same cells before (blue symbols) and after (yellow symbols) naloxone administration in virgin rats and on days 16 and 21 of pregnancy. Note that responses to CCK are increased in pregnancy, but on day 21 there is an opioid suppression of the input. See (155) for details.

For Peer Review Only




Figure 1. The osmosensiveness of magnocellular neurones.

A. Response of a vasopressin cell of the rat supraoptic nucleus to systemic osmotic stimulation in a urethane-anaesthetised rat. The graph plots the firing rate in 1-min bins during continuous i.v. infusion of 1 M NaCl at 26 $\mu\text{l}/\text{min}$ for more than 2 h. The protocol was similar to that in Leng et al. (39), but a lower concentration of NaCl was used over a longer time. The data are from unpublished experiments with Nancy Sabatier used to support the development of a computational model (40). The extracts of spike activity in B-D are from at the beginning of infusion (B); after 4000 s (C); and after 8000 s (D). They show the evolution of intense phasic activity in C, and then of continuous fast spiking activity in D. The minute-by-minute variability of firing rate reflects the intermittent phasic firing pattern, but note the linear rise in mean rate over the duration of infusion.

E. The osmosensiveness of magnocellular neurones involves an increase in afferent input and a direct osmosensitive mechanism. Here in a simple simulation, a fluctuating membrane potential around a mean level 9 mV below the spike threshold (indicated by the red lines) is mimicked by a randomly generated sequence of simplified EPSPs and IPSPs. These are given an amplitude of 3 mV and a half-life of 5 ms, and

arrive at an equal mean rate of 100 Hz for 5 s. None of the fluctuations cross the spike threshold. If the spike threshold is reduced by 2 mV (green line) mimicking the effects of a 2 mV direct depolarization, just one of the fluctuations crosses the threshold (green asterisk). Thus at this level of synaptic input, a small direct depolarization has little effect on spiking activity.

F, In this case, the mean input rate for both EPSPs and IPSPs is 200 Hz. Now, the fluctuations exceed the spike threshold on three occasions (red asterisks), and a depolarization of 2 mV results in an additional 11 threshold crossings (green asterisks). Thus in the presence of sufficient synaptic input, small levels of direct depolarisation can have a large effect on the spiking activity.

190x338mm (96 x 96 DPI)




Figure 2. GABA inputs to vasopressin cells.

In these experiments, single vasopressin cells were recorded from the supraoptic nucleus of a urethane-anaesthetised rat, and single electrical stimuli were applied to the OVLT every 5 s.

A. The experimental set-up. The supraoptic nucleus was exposed by ventral surgery, a stimulating electrode was placed on the neural stalk to allow supraoptic neurons to be antidromically identified, a microdialysis loop was placed on the ventral surface of the nucleus to allow direct application of the GABA antagonist bicuculline, and a stimulating electrode was placed on the OVLT.

B shows interspike interval (ISI) distributions compiled over 1200 s of activity before (orange) and after (blue) bicuculline, showing the resultant increase in mean activity.

C shows mean spike activity before (above) and below (during) bicuculline.

D shows the response to OVLT stimulation before (orange) and after (blue) bicuculline as post-stimulus time histograms (in 1-ms bins) constructed over the periods shown in D. In this cell, OVLT stimulation produced a marked inhibition; bicuculline blocked this inhibition and unmasked an excitatory response. See (74) for details.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

E shows the difference between the two histograms in D, showing the inferred inhibitory effects of OVLT stimulation by subtracting the excitatory effects (blue in D) from the mixed effects (orange in D), indicating that in this cell, the inhibitory effects of OVLT stimulation have a latency of onset of about 20-30 ms.

190x338mm (96 x 96 DPI)




Figure 3. The main regulators of osmosensitiveness in oxytocin cells.

In the rat, extracellular osmotic pressure and increased $[Na^+]$ are sensed both by magnocellular neurones and by cells in the subfornical organ (SFO) and OVLT. Cells in the SFO and OVLT are also responsive to changes in the circulating levels of a number of blood-borne hormones that have important effects on fluid and electrolyte homeostasis. Cells in the OVLT and SFO project directly to the magnocellular neurones, and this direct projection involves the excitatory transmitter glutamate and various peptides. They also project indirectly via the nucleus medianus, and this projection involves the inhibitory transmitter GABA. Astrocytes in the supraoptic nucleus release taurine in response to hypotonic stimulation, and this inhibits via action at glycine receptors on the supraoptic neurones. The GABA-ergic inputs are amplified by nitric oxide, produced by oxytocin cells in an activity-dependent manner (80), and the glutamatergic inputs are moderated by endocannabinoids, also produced by oxytocin cells in an activity-dependent manner (81). Oxytocin release is autoregulated by dynorphin at the level of the nerve terminals in the pituitary. In pregnancy, allopregnanolone enhances the inhibitory effects of GABA, and there is upregulation of both dynorphin expression and down regulation of nitric oxide synthase activity (82).

254x190mm (96 x 96 DPI)




Figure 4. Oxytocin cell responses in pregnancy.

A. In pregnancy, endogenous opioids suppress oxytocin secretion. In these experiments we blocked all opioid actions by pretreating the rats with naloxone to assess opioid-independent changes in the responsiveness of the oxytocin system. Responses were measured in virgin (black bars) and late pregnant rats (day 21, blue bars). The bars show increases in plasma oxytocin concentration (S.E.M.) above basal levels in response to different stimuli. Changes were measured in anaesthetized rats after electrical stimulation of the AV3V region; hypertonic saline, CCK; and in conscious rats following IL-1 β . Both AV3V stimulation and hypertonic saline stimulated oxytocin secretion less effectively in late pregnant rats. Conversely, oxytocin secretory responses to CCK and IL-1 β are greater in late pregnant rats. See (142) for details.

B. Responses of oxytocin cells to CCK (firing rate in 1-min bins above mean basal level) were measured in the same cells before and after naloxone administration in virgin rats and on days 16 and 21 of pregnancy. Note that responses to CCK are increased in pregnancy, but on day 21 there is an opioid suppression of the input. See (148) for details.

254x190mm (96 x 96 DPI)